A convenient green method to synthesize β -carotene from edible carrot and nanoparticle formation

Baker A. Joda

Department of Chemistry, Faculty of Science, University of Kerbala, Karbala, Iraq, baker.judah@uokerbala.edu.iq

Zena M. Abed Al-Kadhim

Department of Chemistry, Faculty of Science, University of Kerbala, Karbala, Iraq

Hussian J. Ahmed

Department of Chemistry, Faculty of Science, University of Kerbala, Karbala, Iraq

Alaa K. H. Al-Khalaf

Department of Environment, College of Environmental Sciences, University of Al-Qasim Green, Iraq

Follow this and additional works at: <https://kijoms.uokerbala.edu.iq/home>

Part of the [Biology Commons](#), [Chemistry Commons](#), [Computer Sciences Commons](#), and the [Physics Commons](#)

Recommended Citation

Joda, Baker A.; Abed Al-Kadhim, Zena M.; Ahmed, Hussian J.; and Al-Khalaf, Alaa K. H. (2022) "A convenient green method to synthesize β -carotene from edible carrot and nanoparticle formation," *Karbala International Journal of Modern Science*: Vol. 8 : Iss. 1 , Article 2.

Available at: <https://doi.org/10.33640/2405-609X.3200>

This Research Paper is brought to you for free and open access by Karbala International Journal of Modern Science. It has been accepted for inclusion in Karbala International Journal of Modern Science by an authorized editor of Karbala International Journal of Modern Science.

University of
Kerbala

A convenient green method to synthesize β -carotene from edible carrot and nanoparticle formation

Abstract

A simple green chemistry approach was used to synthesize β -carotene from edible carrot without using solvent and heat. In this study, beta-carotene was extracted from edible carrots through a natural extraction method with several physical technical steps. The extracted sample was isolated using high-performance liquid chromatography (HPLC). A β -carotene peak was detected from HPLC with an absorbance maximum (2.135 mAU). The absorption of UV-Visible spectrum indicated that beta-carotene absorbs most strongly between 230-300 and 400-500 nm. The functional groups, namely, C=C (1649 cm^{-1}), =C-H ($900\text{-}900\text{ cm}^{-1}$) vibrations of antisymmetric deformation of CH_3 groups and CH_2 groups ($1350\text{-}1450\text{ cm}^{-1}$) of beta carotene were identified by using the FTIR test. The 537, 538, and 539 peaks for all carbon atoms (C-12), one of the 40 carbon atoms is (C-13) and two carbon atoms as (C-13) were indicated using FTMS. The resonance relates to the protons connected to methyl and methylene groups and those attributed to the vinylic protons ($\text{H-C}=\text{C}$) were observed between 1-2 and 5-6 ppm, respectively. The AFM images demonstrate that the structure of the beta-carotene is spherical, and the average diameter of this molecule is 37.22 nm. The findings of the analysis show that the extraction and isolation of β -carotene from the carrot extract was achieved.

Keywords

Green Synthesis, Nano β -Carotene, Medicine Transport Systems, Biomedical Applications, Controlled Drug Release, Pharmaceutical Carriers

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Cover Page Footnote

This research was supported by different Universities such as Kerbala, Al-Qasim Green in Iraq, and New South Wales in Australia.

RESEARCH ARTICLE

A Convenient Green Method to Synthesize β -Carotene from Edible Carrot and Nanoparticle Formation

Baker A. Joda ^{a,*}, Zena M. Abed Al-Kadhim ^a, Hussian J. Ahmed ^a, Alaa K.H. Al-Khalaf ^b

^a Department of Chemistry, Faculty of Science, University of Kerbala, Karbala, Iraq

^b Department of Environment, College of Environmental Sciences, University of Al-Qasim Green, Iraq

Abstract

A simple green chemistry approach was used to synthesize β -carotene from edible carrot without using solvent and heat. In this study, beta-carotene was extracted from edible carrots through a natural extraction method with several physical technical steps. The extracted sample was isolated using high-performance liquid chromatography (HPLC). A β -carotene peak was detected from HPLC with an absorbance maximum (2.135 mAU). The absorption of UV-Visible spectrum indicated that beta-carotene absorbs most strongly between 230-300 and 400-500 nm. The functional groups, namely, C=C (1649 cm^{-1}), =C-H ($900\text{--}900\text{ cm}^{-1}$) vibrations of antisymmetric deformation of CH_3 groups and CH_2 groups ($1350\text{--}1450\text{ cm}^{-1}$) of beta carotene were identified by using the FTIR test. The 537, 538, and 539 peaks for all carbon atoms (C-12), one of the 40 carbon atoms is (C-13) and two carbon atoms as (C-13) were indicated using FTMS. The resonance relates to the protons connected to methyl and methylene groups and those attributed to the vinylic protons (H-C=C) were observed between 1-2 and 5-6 ppm, respectively. The AFM images demonstrate that the structure of the beta-carotene is spherical, and the average diameter of this molecule is 37.22 nm. The findings of the analysis show that the extraction and isolation of β -carotene from the carrot extract was achieved.

Keywords: β -Carotene, Carrot, Nanocarriers, Controlled drug release

1. Introduction

In the recent century, the applications of renewable resources turn out to be more essential for human daily life because they are friendly to nature [1]. The applications and development of green chemistry in academia and industry have been discussed by researchers in the literature [2-5]. The study of nano-sciences especially natural nanomaterials has received increasing attention from many scientific researchers for the progressing trend of nanomaterial applications [6-8]. Generally, edible carrots include the valuable type of carotenoids which are fat-soluble compounds [9]. Carotenoids are colored red-orange pigment that has large, conjugated hydrocarbon skeletons [10]. The basis for the special properties of carotenoids is the conjugated double bond system, which makes them easy to interact with other material due to their delocalized π -electrons having high energy and

needing slight energy for excitation [11]. β -carotene is considered a major of carotenoids (Fig. 1), which is known to show provitamin A activity [10,12]. In addition, β -carotene has antioxidative properties [13,14], therefore, they are expected to reduce the risk of different diseases [15,16]. The consumption of these compounds in different fruits and vegetables ultimately has led to decreased risk to develop such as cancer disease [17-20], cardiovascular diseases [21,22], and those age-related ones [23-25]. In contrast, there are several studies in recent years were found to be contradictory [26,27] and some of these show a negative correlation between β -carotene and some diseases, for example, lung cancer and cardiovascular disease for individuals with smoking activity [28,29].

The possibility of preparing organic molecules using the basis of biomass has previously been investigated. For example, beta-carotene molecules were prepared from the apricot plant as water-

Received 30 April 2021; revised 2 November 2021; accepted 4 November 2021.
Available online 1 February 2022

* Corresponding author at:
E-mail address: baker.judah@uokerbala.edu.iq (B.A. Joda).

<https://doi.org/10.33640/2405-609X.3200>

2405-609X/© 2022 University of Kerbala. This is an open access article under the CC-BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Fig. 1. Molecular structure of beta-carotene [30].

soluble particles using the technique of separating the aqueous phase from the organic layer and diagnosing them as nanoscale particles [31]. The nanoparticles prepared in this way were distinguished by high thermal stability, with nanoscale sizes ranging between (15–250), which were coated with gelatin to be applied in the food and pharmaceutical industry [32]. The effect of nano-encapsulated using β -carotene was investigated by several researchers [33,34]. Recent studies have investigated the possibility of applying nanoencapsulation using beta-carotene molecules in the food sector [35,36]. In addition, the nanoparticles with drugs such as aspirin and metronidazole were applied to verify their ability to nanoencapsulation with high efficiency [37]. The different extraction methods using various types of supplements were utilized and confirmed depending upon the levels of β carotene [38]. This study aims to prepare the nanoparticles of β -carotene from edible carrots by using a convenient green method to prove their ability to nanoencapsulation with high efficiency in part II of the study. Interestingly, there is no study has been previously reported to prepare β -carotene from Iraqi carrots. This synthesizes is important to recognize the nutritional and therapeutic significance of beta carotene.

2. Materials and methods

2.1. β -carotene extraction

The used carrots were obtained from the local supermarket in Karbala, Iraq. The samples were washed using deionized water and cut into tiny pieces to obtain a homogenous sample. The extracted juice was prepared by using a juice extractor without using solvent and heat which may affect the chemical structure of the extracted material or lose the protons that are their presence has a role in the required physical and chemical adsorption and interactions. The extracted sample was centrifuged (15 min) at 7500 rpm and filtered using a piece of cloth. From 100 g of sample, 10 g were selected and kept at -4°C temperature for extraction. The prepared sample (10 g) was homogenized in 30 ml of acetone. The product solution was filtered using Buchner's funnel, then acetone was used to wash the

residue until become colorless. The filtrate was mixed with 20 gm of anhydrous sodium sulfate while the residue was discarded. Then, anhydrous sodium sulfate was separated through filtration and a rotatory evaporator was used to reduce the volume of extract. The extract solution was moved into a volumetric flask (100 ml) and filled with enough acetone and water; therefore, the final extract contains 80% of acetone [36]. Samples of beta carotene (20 μl) were injected into the HPLC system installed at the Laboratory of Chemistry Department, Kerbala University. The chromatographic conditions were Shimadzu HPLC program comprising LC-1000 pump having C18 column connected with LC 250 UV/VIS detector. This instrument was calibrated by using mobile phase (Acetonitrile, dichloromethane, and methanol by the ratio of 70:20:10, respectively) at the rate of 2 ml per minute. Wavelength has been selected at 450 nm, and the column pressure was kept at 1800–2000 PSI.

2.2. Instrumentations

The instruments used in this study were HPLC systems (Shimadzu) containing LC-, and C18 columns connected with UV/Vis detector; Fourier-transform infrared spectroscopy (FTIR) (Shimadzu-8000); Fourier Transform Mass Spectrometry (FTMS); Ultraviolet–Visible (UV–Vis) (Shimadzu) spectrometry, a Bruker 300 MHz NMR Spectrometry and atomic force microscopy (AFM) through Scanning Probe Microscopy (SPM) technique (AA3000). The analyses of FTMS and ^1H NMR were carried out in Australia at the University of New South Wales.

3. Results and discussion

3.1. Separation method

The extract carrot juice was analyzed using HPLC to determine the content of beta carotene in this juice [39]. The results show that the 100 gm of carrot includes thousands of micrograms of beta carotene (11,620 $\mu\text{g}/100\text{ g}$). The findings in this work are approximately in agreement with those described by other studies in which β -carotene quantity in carrot was 11,210 $\mu\text{g}/100\text{ g}$. This variation in beta carotene content is probably due to the influence of several factors such as varieties difference, diameter, extraction methods, column materials, experimental conditions, and used mobile phase [36]. Interestingly, the parameters such as temperature during the analytical process, extraction process of the samples, and the storage time of carrot must be

Fig. 2. UV–Visible spectrophotometry of beta-carotene.

kept in control because they have a tremendous effect on the results [40,41]. A chromatogram chart shows that the maximum peak of β -carotene was identified between 2 and 2.5 min. In addition, there are lower peaks were observed at various wavelengths that may be returned to impurities [42]. The level of purity was determined by measuring the melting point of extracted beta carotene, then compared with the value of pure material. The value of melting point for extracted beta carotene (170°C) was lower than pure material (180°C). This is due to the extracted material including soluble impurities, which are dissolved at lower melting points when compared with pure substance [43].

3.2. UV–visible analysis

Analysis of UV–visible spectrum for beta-carotene is presented in Fig. 2. The instrument conditions (wavelength range: 190–1100 nm, scan mode: single, scan speed: fast, sampling interval: 1, slid

width: 1 nm, S/R exchange: normal) have been used in this analysis. An electronic transition was observed at 210–230 and 400–500 which are revealed to $\pi \rightarrow \pi^*$ transitions because of the electronic conjugation system at the axis of a molecule by double ($\text{C}=\text{C}$) bonds [44]. The main reason is that the energy gap between HOMO and LUMO for conjugate double bonds is less than the energy difference for an isolated double bond. In addition, the λ_{max} increases when the number of conjugated double-bond increases [45].

3.3. FTIR analysis

The dried powder of β -carotene obtained through extracted juice was subjected to FTIR spectroscopy measurements using Shimadzu- 8000 instruments. The sample was combined with KBr, in the sample: KBr ratio of 1:10, then crushed into powder and compressed in a thin layer by a tablet machine at 1 cm^{-1} resolution within the frequency ranges between 4000 and 400 cm^{-1} [37,39]. Beta-carotene is considered as asymmetric conjugated polyenes, so two bands for ($\text{C}=\text{C}$) stretching have appeared, the first around 1649 cm^{-1} assigned to ($\text{C}=\text{C}$) that clear from (CH_3), while the second at 1556 cm^{-1} for ($\text{C}=\text{C}$) bonded with (CH_3) [46,47]. The band at 1456 cm^{-1} is due to scissoring (CH_2) bending, while bands around 1421 cm^{-1} and 1367 cm^{-1} are attributed to asymmetrical and symmetrical (CH_3) bending [48–51]. The oscillations of the $-\text{OH}$ groups of water were found between 3000 and 3600 cm^{-1} as a broad peak [52]. The results of the FTIR spectroscopy study revealed that β -carotene is present in the carrot, as shown in Fig. 3.

Fig. 3. FTIR for the beta-carotene molecule.

3.4. FTMS analysis

Fourier Transform Mass Spectrometry (FTMS $m/z + p$ NSI Full ms [100.00–2000.00]; RT:0.02–0.52AV:10NL:2.26E7; AV: 10 T). In this technique, the cyclotron frequency of ions in a fixed magnetic field is used to determine the m/z . The system provides the highest mass resolution achievable of all mass spectrometers and can provide mass resolutions of 1,000,000 [52–54]. Mass spectroscopy analyzes demonstrated the peaks of β -carotene, as shown in Fig. 4. The beta-carotene nanoparticles have taken the following molar

masses 537.44548, 538.44888, 539.45228, respectively, which are simply corresponded to the molecular ion with a proton added, which is a function of the ionization method used. The 537.44,548 peak is with all carbon atoms are C-12, 538 is with one of the 40 carbon atoms is C-13 (since C-13 is about 1.1% of natural carbon, and the peak at 539 has two carbon atoms as C-13. The resulting compound and molar mass (537.44548 g/mol) belong to the molecular formula ($C_{40}H_{56} + H$), which are formulas with charges in which the proton exchanges by gaining or losing on the surface of the molecule structure which belonging to the beta minutes.

Fig. 4. FTMS spectrum of prepared particles of beta-carotene.

3.5. ¹H NMR spectra

¹H NMR was measured in a Bruker 300 MHz instrument. The ¹H NMR spectra of beta carotene (BC) and carrot extract (CE) are shown in Fig. 5. In comparison between the two spectra, the broad multiple peaks (between 3 and 4 ppm) are found in CE which suggest the presence of a mixture of saccharides [42], while these peaks are disappeared in BC. The resonance at 5 and 6 ppm is attributed to the protons on a double bond, whereas the resonance of protons attached to methyl is observed between 1 and 2 ppm [54]. In addition, the ¹H NMR

Table 1. The properties of surface topology and nanoscale for beta carotene from AFM studies.

Parameter	Beta-Carotene
Core roughness depth (nm)	17.60
Roughness average (nm)	6.89
Surface area ratio	10.9
Average diameter (nm)	37.23
Average height (nm)	27.87

spectra of BC showed the resonances that correspond to the protons attached to the ring were observed between 1.5 and 2.5 ppm, but the aromatic protons are found at 8.2 ppm in CE and disappeared

Fig. 5. ¹H NMR spectrum of (A) carrot extract [42] and (B) Beta Carotene in CDCl₃ at 600 MHz.

Fig. 6. AFM images of Beta-Carotene.

in BC [42]. In conclusion, the FTMS, ¹H NMR, IR, and UV/visible spectra confirmed that the isolated material was beta-carotene.

3.6. AFM analysis

The atomic force microscopy (AFM) is a dynamically evolving field and provides new imaging types to make the technique even more useful, faster, less damaging, and quantitative for polymer science applications [55]. AFM/SPM mapping was performed to evaluate the topography, surface potential distribution, and nanoscale properties for the beta-carotene molecules [56]. Table 1 and Fig. 6 show the values of core roughness depth (17.60 nm), surface roughness average (6.89 nm), and surface area ratio (10.9), along with a smaller average of diameter (37.23 nm) and height (27.87 nm). These properties have many useful advantages, therefore can be used in different vital aspects such as medicine and industrial applications such as pharmaceutical carriers.

4. Conclusions

In this work, natural extraction along with several physical technical steps was used to prepare a beta carotene by using an edible carrot. The extracted beta carotene has been characterized using HPLC, UV–Vis, FTIR, FTMS, and ¹H NMR spectroscopy. These techniques confirmed that the isolated material was beta-carotene. The surface and nanoscale properties were investigated using AFM. The properties are confirmed that beta-carotene has many useful advantages, which can use in different medicine and industrial applications.

Acknowledgment

This research was supported by different Universities such as Kerbala, Al-Qasim Green in Iraq, and New South Wales in Australia.

References

- [1] P. Phanthong, P. Reubroycharoen, X. Hao, G. Xu, A. Abudula, G. Guan, Nanocellulose: extraction and application, *Carbon Resour. Convers.* 1 (2018) 32–43, <https://doi.org/10.1016/j.crcon.2018.05.004>.
- [2] P.M. Nowak, R. Wietecha-Postuszny, J. Pawliszyn, White analytical chemistry: an approach to reconcile the principles of green analytical chemistry and functionality, *Trac. Trends Anal. Chem.* 138 (2021) 116223, <https://doi.org/10.1016/j.trac.2021.116223>.
- [3] T.F. Alessandr, T.W. Cabri, A translation of the twelve principles of green chemistry to guide the development of cross-coupling reactions, *Catal. Today*, Available online 17 September (2021), (In Press). <https://doi.org/10.1016/j.cattod.2021.09.022>.
- [4] B.L. Eilks, A systematic review of the green and sustainable chemistry education research literature in mainland China, *Sustainable Chem. Pharm.* 21 (2021) 100446, <https://doi.org/10.1016/j.scp.2021.100446>.
- [5] D. Lenoir, K.W. Schramm, J.O. Lalah, Green Chemistry: some important forerunners and current issues, *Sustainable Chem. Pharm.* 18 (2020) 100313, <https://doi.org/10.1016/j.scp.2020.100313>.
- [6] M.E.T. Yazdi, Mohammad Sadegh Amiri, Setareh Akbari, Mohammad Sharifalhoseini, F. Nourbakhsh, M. Mashreghi, et al., Green synthesis of silver nanoparticles using *Helichrysum graveolens* for biomedical applications and wastewater treatment, *Bio. Nano Sci.* 10 (2020) 1121–1127, <https://doi.org/10.1007/s12668-020-00794-2>.
- [7] G.L.R. Oliveira, I. Medeiros, S.S.d. Nascimento, R.L.S. Viana, D.L. Porto, H.A.O. Rocha, et al., Carotenoid rich-extract from Cantaloupe melon (*Cucumis melo* L.) nanoencapsulated in gelatin under different storage conditions, *Food Chem.* 34 (2021) 129055, <https://doi.org/10.1016/j.foodchem.2021.129055>.
- [8] P.T.T. Phuong, S. Lee, C.h. Lee, B. Seo, S. Park, K. Take Oh, et al., Beta-carotene-bound albumin nanoparticles modified with chlorin e6 for breast tumor ablation based on photodynamic therapy, *Colloids Surf. B Biointerfaces* 171 (2018) 123–133, <https://doi.org/10.1016/j.colsurfb.2018.07.016>.
- [9] D.M. Yebra, S. Kiil, K. Dam-Johansen, Antifouling technology-past, present and future steps towards efficient and environmentally friendly antifouling coatings, *Prog. Org. Coating* 50 (2004) 75–104, <https://doi.org/10.1016/j.porgcoat.2003.06.001>.
- [10] X.D. Wang, Carotenoids, in: C.A. Ross, B. Caballero, R.J. Cousins, K.L. Tucker, T.R. Ziegler (Eds.), *Modern Nutrition in Health and Disease*, eleventh ed., Lippincott Williams & Wilkins, 2014, pp. 427–439.
- [11] J. Clayden, N. Greeves, S. Warren, P. Wothers, *Delocalization, and Conjugation*, Oxford University Press, Oxford, New York, United States, 2009, pp. 156–171.

- [12] J.A. Olsen, Provitamin A function of carotenoids: the conversion of β -carotene into vitamin A, *J. Nutr.* 119 (1989) 105–108, <https://doi.org/10.1093/jn/119.1>.
- [13] W. Stahl W, H. Sies, Carotenoids and flavonoids contribute to nutritional protection against skin damage from sunlight, *Mol. Biotechnol.* 37 (1) (2007) 26–30, <https://doi.org/10.1007/s12033-007-0051-z>.
- [14] S.A.R. Paiva, R.M. Russell, Review series: antioxidants and their clinical applications β -carotene and other carotenoids as antioxidants, *J. Am. Coll. Nutr.* 18 (5) (1999) 426–433, <https://doi.org/10.1080/07315724.1999.10718880>.
- [15] A.T. Diplock, J.L. Charleux, G. Crozier-Willi, F.J. Kok, C. Rice-Evans, M. Roberfroid, et al., Functional food science and defence against reactive oxidative species, *Br. J. Nutr.* 80 (1998) 77–112, <https://doi.org/10.1079/bjn19980106>.
- [16] N.I. Krinsky, The biological properties of carotenoids, *Pure Appl. Chem.* 66 (5) (1994) 1003–1010.
- [17] E. Giovannucci, A. Ascherio, E.B. Rimm, M.J. Stampfer, G.A. Colditz, W.C. Willett, Intake of carotenoids and retino in relation to risk of prostate cancer, *J. Natl. Cancer Inst.* 87 (23) (1995) 1767–1776, <https://doi.org/10.1093/jnci/87.23.1767>.
- [18] A. Braithwaite, F.J. Smith, *Chromatographic Methods*, 4th., Chapman & Hall, London, 1995.
- [19] F.L. Meyskens, A. Manetta, Prevention of cervical intraepithelial neoplasia and cervical cancer, *Am. J. Clin. Nutr.* 62 (6) (1995) 1417–1419, <https://doi.org/10.1576/toag.5.1.21>.
- [20] W. Zheng, W.J. Blot, E.J. Diamond, E.P. Norkus, V. Spate, J.S. Morris, et al., Serum micronutrients and the subsequent risk of oral and pharyngeal cancer, *Cancer Res.* 53 (4) (1993) 795–798.
- [21] J.M. Gaziano, Antioxidant vitamins, and coronary artery disease risk, *Am. J. Med.* 97 (3) (1994) 18–21, [https://doi.org/10.1016/0002-9343\(94\)90294-1](https://doi.org/10.1016/0002-9343(94)90294-1).
- [22] J.M. Gaziano, C.H. Hennekens, The role of beta carotene in presentation of cardiovascular diseases, *Ann. N. Y. Acad. Sci.* 1 (1993) 148–155, <https://doi.org/10.1111/j.1749-6632.1993.tb26166.x>.
- [23] J.M. Seddon, U.A. Ajani, R.D. Sperduto, R. Hiller, N. Blair, T.C. Burton, et al., The eye disease case-control study group. Dietary carotenoids, vitamins A, C, and E, and advanced age-related macular degeneration, *JAMA: J. Am. Med. Assoc.* 272 (18) (1994) 1413–1420, <https://doi.org/10.1001/jama.1994.03520180037032>.
- [24] R.D. Semba, F. Lauretani, L. Ferrucci, Carotenoids as protection against sarcopenia in older adults, *Arch. Biochem. Biophys.* 458 (2) (2007) 141–145, <https://doi.org/10.1016/j.jabb.2006.11.025>.
- [25] S.T. Mayne, Beta-carotene, carotenoids, and disease prevention in humans, *Faseb. J.* 10 (7) (1996) 690–701, <https://doi.org/10.1096/fasebj.10.7.8635686>.
- [26] R. Greenberg, J.A. Baron, T.D. Tosteson, D.H. Freeman, G.J. Beck, J.H. Bond, et al., for the Polyp Prevention Study Group, A clinical trial of antioxidant vitamins to prevent colorectal adenoma, *N. Engl. J. Med.* 331 (1994) 141–147, <https://doi.org/10.1056/NEJM199407213310301>.
- [27] M. Mozaffarieh, S. Sacu, A. Wiedrich, The role of the carotenoids, lutein, zeaxanthin, in protecting against age-related macular degeneration: a review based on controversial evidence, *Nutr. J.* 2 (20) (2003) 1–27, <https://doi.org/10.1186/1475-2891-2-20>.
- [28] O.P. Heinonen, D. Albanes, The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers, *N. Engl. J. Med.* 330 (15) (1994) 1029–1035, <https://doi.org/10.1056/NEJM199404143301501>.
- [29] G.S. Omenn, G.E. Goodman, M.D. Thornquist, J. Balmes, M.R. Cullen, A. Glass, et al., Risk factors for lung cancer and intervention effects in CARET, the beta-carotene and retinol efficacy trial JNCI, *J. Natl. Cancer Inst.* 88 (21) (1996) 1550–1559, <https://doi.org/10.1093/jnci/88.21.1550>.
- [30] P. Atkins, J. de Paula (Eds.), *Physical Chemistry*, eighth ed., Oxford Press, Oxford, UK, 2006.
- [31] D. Horn, E. Luddecke, Preparation and characterization of nano-sized carotenoid hydrosols, in: E. Pelizzani (Ed.), *Fine Particles Science and Technology*, Kluwer Academic Publishers, 1996, pp. 761–775, <https://doi.org/10.1002/app.43027>.
- [32] J. Kim, T. Seo, S. Lim, Preparation of an aqueous dispersion of β -carotene nanocomposites through complex formation with starch dextrin, *Food Hydrocolloids* 33 (2013) 256–263, <https://doi.org/10.1016/j.foodhyd.2013.04.001>.
- [33] G. Liu, Y. Zhou, L. Chen, Intestinal uptake of barley protein-based nanoparticles for β -carotene delivery, *Acta Pharm. Sin. B* 9 (1) (2019) 87–96, <https://doi.org/10.1016/j.apsb.2018.10.002>.
- [34] M. Battistonia, R. Bacchettaa, F. Di Renzoa, F. Metrucciob, E. Menegola, Effect of nano-encapsulation of β -carotene on *Xenopus laevis* embryos development (FETAX), *Toxicol. Rep.* 7 (2020) 510–519, <https://doi.org/10.1016/j.toxrep.2020.04.004>.
- [35] F.J. Gutierrez, S. Albillos, E. Casas-Sanz, M. Mussons, Methods for the nanoencapsulation of β -carotene in the food sector, *Trends Food Sci. Technol.* 32 (2013) 73–83, <https://doi.org/10.1016/j.tifs.2013.05.007>.
- [36] M. Nauman Ahamad, M. Saleemullah, Hamid Ullah Shah, Iqtidar A. Khalil, A.U.R. Saljoqi, Determination of beta carotene content in fresh vegetables using HPLC, *Sarhad J. Agric.* 23 (3) (2007) 767–770.
- [37] Z.M. Abed Al-Kadhim, B.A. Joda, A.K.H. Al-Khalaf, A convenient green method to synthesize a nanocellulose from edible fresh potato, *J Conf Proc Conf 9th* (in press).
- [38] N. Hazuki, Extraction of Carotenoids from Natural Products and Nanoparticle Formation Using Supercritical Fluid, Ph.D. Thesis, Nagoya University, 2015, pp. 1–114.
- [39] C.h. Zhang, Y. Fu, Z. Li, T. Li, Y. Shi, H. Xie, et al., Application of whey protein isolate fibrils in encapsulation and protection of β -carotene, *Food Chem.* 346 (2021) 128963, <https://doi.org/10.1016/j.foodchem.2020.128963>.
- [40] C. Marcela, R.B.D. Amaya, Carotenoid composition of cooked green vegetables from restaurants, *J. Food Chem.* 83 (2003) 595–600.
- [41] I.A. Khalil, F.R. Varanani, Carotinioid extraction and analysis by reversed-phase HPLC system, *Sarhad J. Agric.* 105 (67) (1996) 15–21.
- [42] M. Jayanthi, S. Megarajan, S. Bala, R.K. Kamlekar, A. Veerappan, A convenient green method to synthesize luminescent carbon dots from edible carrot and its application in bioimaging and preparation of nanocatalyst, *J. Mol. Liq.* 278 (2019) 175–182, <https://doi.org/10.1016/j.jmolliq.2019.01.070>.
- [43] Steven A. Hardinger, A simple demonstration of the effect of impurities on melting point, *J. Chem. Edu.* Easton 72 (3) (1995) 191–286, <https://doi.org/10.1021/ed072p250>.
- [44] Yu Liu, Philip Kilby, J.F. Terry, W. Timothy, Schmidt, Electronic transitions of molecules: vibrating Lewis structures, *Chem. Sci.* 10 (2019) 6809–6814, <https://doi.org/10.1039/C9SC02534K>.
- [45] J. Wang, Q. Liu, B. Xie, Z. Sun, Effect of ultrasound combined with ultraviolet treatment on microbial inactivation and quality properties of mango juice, *Ultrason. Sonochem.* 64 (2020) 105000, <https://doi.org/10.1016/j.ultsonch.2020.105000>.
- [46] S. Lotfy, Y.H.A. Fawzy, Characterization and enhancement of the electrical performance of radiation modified poly (vinyl) alcohol/gelatin copolymer films doped with carotene, *J. Rad. Res. Appl. Sci.* 7 (2014) 338–345, <https://doi.org/10.1016/j.jrras.2014.04.003>.
- [47] T. Neha, T. Shishir, D. Ashutosh, Fourier transform infrared spectroscopy (FTIR) profiling of red pigment produced by *Bacillus subtilis* PD5, *Afr. J. Biotechnol.* 16 (27) (2017) 1507–1512, <https://doi.org/10.5897/AJB2017.15959>.
- [48] M. Kreck, P. Kuerbel, M. Ludwig, P.J. Paschold, H. Dietrich, Identification and quantification of carotenoids in pumpkin cultivars (*Cucurbita maxima* L.) and their juices by liquid chromatography with ultraviolet-diode array detection, *J. Appl. Bot. Food Qual.* 80 (2006) 93–99.
- [49] K.V. Berezin, V.V. Nechaev, Calculation of the IR spectrum and the molecular structure of β -carotene, *J. Appl.*

- Spectrosc. 72 (2005) 164–171, <https://doi.org/10.1007/s10812-005-0049-x>.
- [50] S. Schlücker, A. Szeghalmi, M. Schmitt, J. Popp, W. Kiefer, Density functional and vibrational spectroscopic analysis of β -carotene, *J. Raman Spectrosc.* 34 (2003) 413–419, <https://doi.org/10.1002/jrs.1013>.
- [51] N. Quijano-Ortega, C.A. Fuenmayor, C.Z. Zuluaga-Dominguez, C.D. Diaz-Moreno, S. Ortiz-Grisales, M. García-Mahecha, et al., FTIR-ATR Spectroscopy combined with multivariate regression modeling as a preliminary approach for carotenoids determination in cucurbita spp, *Appl. Sci.* 10 (2020) 3722, <https://doi.org/10.3390/app10113722>.
- [52] G.A. Marshall, L. C.h. Hendrickson, S.G. Jackson, Fourier transform ion cyclotron resonance mass spectrometry: a primer, *Mass Spectrom. Rev.* 17 (1) (1998) 1–35.
- [53] Marcus Höring, Christer S. Ejsing, Sabrina Krautbauer, Verena M. Ertl, Ralph Burkhardt, Gerhard Liebisch, Accurate quantification of lipid species affected by isobaric overlap in Fourier-transform mass spectrometry, *J. Lipid Res.* 62 (2021) 100050, <https://doi.org/10.1016/j.jlr.2021.100050>.
- [54] M.H. Wang, R.Y. Wang, X.Y. Wei, W. Zhao, X. Fan, Molecular characteristics of the oxidation products of a lignite based on the big data obtained from Fourier transform ion cyclotron resonance mass spectrometry, *Fuel* 295 (1) (2021) 120644, <https://doi.org/10.1016/j.fuel.2021.120644>.
- [55] Y. Liu, G.J. Vancso, Polymer single chain imaging, molecular forces, and nanoscale processes by Atomic Force Microscopy: the ultimate proof of the macromolecular hypothesis, *Prog. Polym. Sci.* 104 (2020) 101232, <https://doi.org/10.1016/j.progpolymsci.2020.101232>.
- [56] E. Rahimi, R. Offoiaich, S. Hosseinpour, A. Davoodi, K. Baert, A. Lutz, et al., Effect of hydrogen peroxide on bovine serum albumin adsorption on Ti6Al4V alloy: a scanning Kelvin probe force microscopy study, *Appl. Surf. Sci.* 563 (15) (2021) 150364, <https://doi.org/10.1016/j.apsusc.2021.150364>.