Molecular characterization of wild *Pleurotus ostreatus* (MW457626) and evaluation of β -glucans polysaccharide activities

Ghazwan Q. Hasan Ph.D.

Department of Biology, College of Education for Pure Sciences, University of Mosul, Iraq, ghazwan.qasim@yahoo.com

Shimal Y. Abdulhadi Ph.D.

Department of Biology, College of Education for Pure Sciences, University of Mosul, Iraq

Follow this and additional works at: <https://kijoms.uokerbala.edu.iq/home>

Part of the [Biology Commons](#), [Chemistry Commons](#), [Computer Sciences Commons](#), and the [Physics Commons](#)

Recommended Citation

Hasan, Ghazwan Q. Ph.D. and Abdulhadi, Shimal Y. Ph.D. (2022) "Molecular characterization of wild *Pleurotus ostreatus* (MW457626) and evaluation of β -glucans polysaccharide activities," *Karbala International Journal of Modern Science*: Vol. 8 : Iss. 1 , Article 6.

Available at: <https://doi.org/10.33640/2405-609X.3204>

This Research Paper is brought to you for free and open access by Karbala International Journal of Modern Science. It has been accepted for inclusion in Karbala International Journal of Modern Science by an authorized editor of Karbala International Journal of Modern Science.

Molecular characterization of wild *Pleurotus ostreatus* (MW457626) and evaluation of β -glucans polysaccharide activities

Abstract

Pleurotus ostreatus is a common cultivated edible mushroom worldwide. The fruiting bodies of *P. ostreatus* is a rich source of a β -glucans polysaccharide. The current study aimed to investigate the effectiveness of β -glucans as a natural polysaccharide produced by *P. ostreatus* as an antioxidant, antimicrobial, and anticancer. The molecular identification of *P. ostreatus* isolate was confirmed by Internal Transcribed Spacer (ITS) sequence. The sequence alignment and phylogenetic evolutionary relationship of studied ITS sequence were performed against some deposited sequences in GenBank. The analysis of high-performance liquid chromatography (HPLC) as well as the result of fourier transform infrared spectroscopy (FTIR) has confirmed the presence of β -glucans polysaccharide in the tested samples. The percentage of antioxidant activity of β -glucans showed a gradual increase from 8.59% to 12.36, 18.56, 23.69, 44.66 and 80.36% at the concentrations of 31.2, 64.4, 125, 250, 500, and 800 μ g/ml, respectively. In addition, all concentrations of β -glucans showed higher antioxidant activities when compared with standard antioxidant (Vitamin C). The highest- antimicrobial activity of β -glucans polysaccharide was against *P. aeruginosa* with a zone of inhibition (45 mm), while the lowest activity was against *S. aureus* (13 mm) both at 100 mg/mL. The percentage of growth-inhibiting of MCF-7 a human breast cancer cell line and normal WRL-68 cell line affected by β -glucans were determined by 3-(4,5)-dimethylthiazol (-z-y1)-3,5-di-phenyltetrazoliumromide (MTT assay). The results revealed that the percentage of growth inhibiting of MCF-7 and WRL-68 cells were gradually increased in both lines and was the highest with MCF-7 line, where the percentage were 18, 23, 50, 59, and 62 % compared with WRL-68 line which showed 4, 6, 9, 13, and 22% at 1.0, 1.5, 2.0, 2.5, and 3.0 μ g/ml, respectively. This present paper revealed that *P. ostreatus* has therapeutic values that can be used as a natural medicine instead of industrial compounds.

Keywords

Pleurotus ostreatus, β -glucans, ITS region, Antioxidant, Antimicrobial, Anticancer, MTT assay

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License](https://creativecommons.org/licenses/by-nc-nd/4.0/).

RESEARCH PAPER

Molecular Characterization of Wild *Pleurotus Ostreatus* (MW457626) and Evaluation of β -glucans Polysaccharide Activities

Ghazwan Q. Hasan*, Shimal Y. Abdulhadi

Department of Biology, College of Education for Pure Sciences, University of Mosul, Iraq

Abstract

Pleurotus ostreatus is a common cultivated edible mushroom worldwide. The fruiting bodies of *P. ostreatus* is a rich source of a β -glucans polysaccharide. The current study aimed to investigate the effectiveness of β -glucans as a natural polysaccharide produced by *P. ostreatus* as an antioxidant, antimicrobial, and anticancer. The molecular identification of *P. ostreatus* isolate was confirmed by Internal Transcribed Spacer (ITS) sequence. The sequence alignment and phylogenetic evolutionary relationship of studied ITS sequence were performed against some deposited sequences in GenBank. The analysis of high-performance liquid chromatography (HPLC) as well as the result of fourier transform infrared spectroscopy (FTIR) has confirmed the presence of β -glucans polysaccharide in the tested samples. The percentage of antioxidant activity of β -glucans showed a gradual increase from 8.59% to 12.36, 18.56, 23.69, 44.66 and 80.36% at the concentrations of 31.2, 64.4, 125, 250, 500, and 800 μ g/ml, respectively. In addition, all concentrations of β -glucans showed higher antioxidant activities when compared with standard antioxidant (Vitamin C). The highest-antimicrobial activity of β -glucans polysaccharide was against *P. aeruginosa* with a zone of inhibition (45 mm), while the lowest activity was against *S. aureus* (13 mm) both at 100 mg/mL. The percentage of growth-inhibiting of MCF-7 a human breast cancer cell line and normal WRL-68 cell line affected by β -glucans were determined by 3-(4,5)-dimethylthiazol (-z-y1)-3,5-di-phenyltetrazoliumromide (MTT assay). The results revealed that the percentage of growth inhibiting of MCF-7 and WRL-68 cells were gradually increased in both lines and was the highest with MCF-7 line, where the percentage were 18, 23, 50, 59, and 62% compared with WRL-68 line which showed 4, 6, 9, 13, and 22% at 1.0, 1.5, 2.0, 2.5, and 3.0 μ g/ml, respectively. This present paper revealed that *P. ostreatus* has therapeutic values that can be used as a natural medicine instead of industrial compounds.

Keywords: *Pleurotus ostreatus*, β -glucans, ITS region, Antioxidant, Antimicrobial, Anticancer, MTT assay

1. Introduction

Mushrooms have been previously played a crucial role in folk medicine [1], and utilized in popular medicine of China and some Asian countries. They have antioxidant, antibacterial anticancer, antiviral, anti-inflammatory, hypoglycemic, and hypocholesterolemic characteristics [2–4]. Nowadays, edible mushrooms are popular beneficial foods with low calories, carbohydrates, fat and cholesterol-free [5]. Mushrooms contain some metal elements and vitamins such as calcium, potassium,

phosphorus, copper, ascorbic acid, vitamin D and K, niacin, thiamine and others [6]. The content of macro and micronutrients in *Pleurotus* spp. varies according to the growth substrate [7]. The content of macro and micronutrients in *Pleurotus* spp. varies according to the growth substrate. For instance, Ca and K content in *Pleurotus* spp. were ranged from 1 to 330, and from 271 to 4054.3 mg/100 g d.w. respectively [7]. Moreover, vitamin D2 content in *Pleurotus* spp is low, but *Pleurotus* spp. have a high content of ergosterol, which generates vitamin D2 upon UV light exposure [8] They are rich also in

Received 19 August 2021; revised 6 November 2021; accepted 10 November 2021.
Available online 11 February 2022

* Corresponding author at:
E-mail addresses: dr.ghazwan@uomosul.edu.iq, ghazwan.qasim@yahoo.com (G.Q. Hasan).

<https://doi.org/10.33640/2405-609X.3204>

2405-609X/© 2022 University of Kerbala. This is an open access article under the CC-BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

some secondary metabolites which have high medicinal value such as polysaccharide, lecithin, phenolic compounds, terpenoids, saponin, polyketides, and others [9,10]. Several studies revealed that the *Pleurotus* genus appears multidirectional benefits for supporting human health [11–13]. Further, the oyster mushroom was considered a functional food, due to its positive role in a human being [14,15]. The *Pleurotus* genus consists of around 14 species [16]. Mushrooms are enriched with minerals, proteins, and vitamins [17]. The therapeutic properties and bioactive compounds were found in both fruiting bodies and the mycelia of oyster mushrooms [18]. There are two different compounds found in *Pleurotus* mushrooms according to their molecular weight, high-molecular weight secondary metabolites which mainly consist of polysaccharides, involving β -glucans, proteins and peptides; and low-molecular weight secondary metabolites involve polyphenols, terpenes and fatty acids [19]. The outstanding compounds in *Pleurotus* mushrooms are the β -glucans polysaccharides which are mainly found in the cell wall of mushrooms. It involves glucopyranose units binding with glycosidic bonds at the positions of (1 $\rightarrow$ 3)- β , (1 $\rightarrow$ 6)- β - or (1 $\rightarrow$ 3)- α . β -glucans. It has many important roles such as activation of the human immune system, antioxidant, antimicrobial, anticancer, antiviral, antifungal, controlling the cholesterol level and glucose regulation in blood [20]. The *G. lucidum* polysaccharides have also similar activity of anticancer [21,22]. β -glucans inhibits the growth of cancer cells by stimulating the immune reaction in the active-cells [23]. It was found that β -glucans encourages the inducing of interferon-gamma and interleukin-12 (IL-12) in lymphocytes [24] and increasing of T helper 1 response through inducing of interleukin-1 (IL-1), nitric oxide and tumor necrosis factor-alpha [25]. Oyster mushroom is one of many richest sources of β -glucans polysaccharide [26]. Based on [27] the total glucans in *P. ostreatus* is 25.636 g/100 g dm (dry mass), while the β -glucans concentration is (24.230 g/100g dm). However, the total concentrations of β -glucose were (4.6 g/100 g dm and 9 g/100 g dm) in mycelia and fruiting bodies of *P. ostreatus*, respectively [28]. FTIR technique helps to detect the active groups that appear in the sample and is commonly used to examine the fungi and plant-based polysaccharides [22,29]. The ITS region is widely considered to be the most important sequenced DNA region used for molecular detection purposes in fungi [30,31] and is considered the universal barcode sequence in fungi [32]. The purposes of the present study are to

perform some molecular analysis such as multiple sequence alignments and draw a phylogenetic tree of studied ITS sequence with deposited sequences in GenBank, and study the antioxidant, antimicrobial, and anticancer activities. The macroscopic features of the Basidiomycota were recorded, based on the approach followed by Ref. [34]. The collected isolates were deposited at the Herbário do Estado Maria Eneyda P. Kaufmann Fidalgo (SP) under the following names CCB001, Pt II, Pt III, and Pt IV and their accession numbers were SP392848, SP392847, SP392849, and SP392850, respectively. The microscopic analysis involved rehydrates the dry material with 70% ethanol, then with 5% KOH reagent [35]. For more morphological characterizations, a consultation was needed from Refs. [36–45].

2. Materials and methods

2.1. Sampling and isolation of *P. ostreatus*

P. ostreatus fruiting bodies growing on the trunks of Morus tree (Fig. 1) have been collected from the farms located in Rashidiyah region/Mosul city, Iraq <https://goo.gl/maps/tFJwGPjRHrPWbUFD8>. Samples were placed in sterile polyethylene bags and transferred to the lab, then washed with running water several times, after which they were cut lengthwise into small parts in thin slices. The pieces were sterilized with 70% ethyl alcohol for 2 min, then washed with sterile distilled water for 2–5 minutes, after which the samples were left to dry on sterile filter papers Whatman No. 1, then the pieces were distributed on Petri dishes containing Potato Dextrose Agar using sterile forceps with alcoholic flame. The dishes were covered with aluminium foil and transferred to the incubator at 28 °C for 7–10 days. Growing mycelia were preserved in a suitable place until use [33].

2.2. Identification

2.2.1. Phenotypic identification

The macroscopic features of the Basidiomycota were recorded, based on the approach followed by Ref. [34]. The collected isolates were deposited at the Herbário do Estado Maria Eneyda P. Kaufmann Fidalgo (SP) under the following names CCB001, Pt II, Pt III, and Pt IV and their accession numbers were SP392848, SP392847, SP392849, and SP392850, respectively. The microscopic analysis involved rehydrates the dry material with 70% ethanol, then with 5% KOH reagent [35]. For more morphological characterizations, a consultation was needed from Refs. [36–45].

2.2.2. Molecular identification

Genomic DNA from pure mycelia of *P. ostreatus* was isolated using the CTAB method as described by Ref. [46]. The DNA fragment of ITS was amplified by PCR using ITS1 forward (5'- TCCGTAGGTGA-ACCTGCGG -3') and ITS4 reverse (5'- TCCTCCG-CTTATTGATATGC-3') primers. The DNA fragment of 657 bp was checked by electrophoresis on a 1% agarose gel and visualized by a UV transilluminator (Fig. 2). The PCR product was sent for sequencing to Microgen Company/South Korea <https://dna.macrogen.com/#>. The ITS sequence analysis of *P. ostreatus* was performed with the BLASTn site at the NCBI <https://blast.ncbi.nlm.nih.gov/Blast.cgi>.

2.3. Extraction of β -glucans

The β -glucans polysaccharide was extracted from *P. ostreatus* using the approach presented by Ref. [47]. The cultured mycelia were extracted by hot water for 60 min, the culture was then filtered with Whatman No. 1 filter paper. The samples were precipitated with 4 vol of 95% (v/v) ethanol and left for a day at 4 °C. Next, the centrifugation of preprecipitates was performed at 10 000×g for 10 min, and the supernatant was discarded. The precipitates were lyophilized into powders. The powders then were redissolved in deionized water, and the final stock concentration was 20 mg/ml, and then kept at 4 °C until use.

2.4. β -glucans activities

2.4.1. Measurement of antioxidant efficacy of β -glucans

The purified β -glucans isolated from *P. ostreatus* were subjected to antioxidant measurement using

2,2-diphenyl-1-picrylhydrazyl or DPPH radical scavenging technique [48]. The stock concentration of 400 μ g/ml was made by dissolving 0.04 g of DPPH into 100 ml of methanol. The control solution was made by dissolving 0.5 g of ascorbic acid into 100 ml of methanol and dH₂O in a 1:1 ratio. The concentrations (31.2, 64.4, 123, 250, 500 and 800 μ g/ml) of β -glucans and ascorbic acid were prepared (Fig. 6). 1 ml of DPPH was mixed with 1 ml of each prepared concentration of β -glucans and ascorbic acid, the mixture then was vigorously shaken and left in dark at room temperature for a half an hour. The control solution contained the same volume as the reaction solution except using methanol solvent instead of the extracts above. The absorbance of DPPH was determined at the wavelength 517 nm. The DPPH value was expressed as a percentage, according to the method used by Ref. [49] as in the following equation:

$$\text{Percentage of scavenging} = A^O - A1 / A^O \times 100$$

A^O = Absorbance of control;

Fig. 1. *Morus* tree carrying fruiting bodies of *P. ostreatus*. Scale bar: 1 cm.

Fig. 2. Agarose gel image of amplified ITS fragment (657 bp) of *P. ostreatus*. (1) 100 bp DNA ladder, (2) ITS fragment amplified with ITS1 forward and ITS4 reverse primers.

Fig. 3. (Continued).

A1 = Absorbance of sample.

2.4.2. Assessment of antimicrobial activity of β -glucans

The antimicrobial properties of β -glucans were accomplished with the agar well diffusion method [50]. The β -glucans extract was dissolved in Dimethyl sulfoxide of 100 mg/mL, the chloramphenicol at 40 mg/mL and dH₂O were used as positive and negative controls, respectively. The selective wells poured into Petri dishes (5-mm in diameter) containing 2% nutrient agar (2% agar, 0.5% Peptone, 0.5% sodium chloride, 0.3% beef extract, 1 L distilled water, at pH 7.0). The test was accomplished in triplicate, and the cultures were kept overnight at 28 °C. For antimicrobial activity, the plates were incubated at convenient growth temperatures for tested bacteria (*Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus*, and *Proteus mirabilis*). The diameter of the inhibition zone was measured in mm [51] as shown in Fig. 7 and Table 3.

2.4.3. Determination of cytotoxic activity of β -glucans

MCF-7 a human breast cancer cell line and WRL-68 a normal cell line were kindly provided from the Biotechnology Research Center of Al-Nahrain University/Baghdad and used to determine the anti-cancer activity of β -glucans (Table 4). The MTT assay is a proliferation detection and colorimetric cytotoxicity test, this approach is depending on the metabolic effects of normal cells [52]. The tetrazolium salts are usually decreased in variable cells to a blue-coloured formazan, which is proportional to the number of active cells. The MCF-7 and WRL-68 cell lines (5×10^5 cells/mL) were placed into a 96-well microplate, the cultures were grown overnight at standard conditions of 5% CO₂ ml at 37 °C. The cells were incubated for 48 hours with different concentrations of β -glucans (1.0, 1.5, 2.0, 2.5, and 3.0 μ g/ml) (Table 4). Each sample was prepared with 1%

of fetal bovine serum (FCS) medium. Following incubation, 1 mg/mL of MTT was added to each sample and incubated once again for 3 h. Then, 100 μ L of the sodium dodecyl sulfate solution were applied to each sample to dissolve formazan crystals, and the wells were incubated overnight at 37 °C. The optical density was determined at 570 nm, and then the cytotoxicity of the cancer cell lines was calculated as a percentage based on (100% of living cells in the control).

3. Results

3.1. Isolation of genomic DNA, PCR amplification, and ITS sequence analysis

Genomic DNA from fruiting bodies of *P. ostreatus* was extracted using the CTAB method. The ITS fragment 657 bp was successfully amplified by PCR using forward (ITS1) and reverse (ITS4) primers and clearly shown on gel electrophoresis (Fig. 2). The PCR product was sent for sequencing to Microgen Company/South Korea <https://dna.macrogen.com/#>. The ITS sequence was collected in high quality and then deposited in GenBank under accession number MW457626 which showed a homology of 99% with some deposited sequences in GenBank (Accessions: MK603976, MK281340, MK281339, MH287458.1 and KT968340.1). Multiple sequence alignments and evolutionary relationships were performed for the ITS region by MEGA X_10.2.2 software <https://www.megasoftware.net/>. (Fig. 3: A and B).

3.2. HPLC analysis of β -glucans polysaccharide

1 mL of extract was analyzed by HPLC [57]. This analysis was accomplished in the Department of Environment and Water/Ministry of Science and Technology, following the acid hydrolysis process using the HPLC device (SYKAM model-Germany). The used column was C18-ODS with dimensions 4.6 mm* 25 cm. The measurements were taken by (UV–Vis) at 360 nm wavelength with a flow rate 1 mL per minute. The analysis revealed the presence of β -glucans polysaccharide as a main bioactive compound (Fig. 4 and Table 1).

3.3. Fourier-transform infrared spectroscopy analysis of β -glucans polysaccharide

FTIR is an assay used to collect an infrared spectrum of absorption or emission from different states of matter (solid, liquid or gas). FTIR spectrum analysis pointed to the existence of β -glycosidic

Fig. 4. HPLC chromatogram of polysaccharide β -glucans extracted from *P. ostreatus*. The samples were detected by (UV–Vis) at a wavelength of 360 nm at a flow rate 1 mL per minute.

Fig. 5. FTIR spectra of β -glucans extracted from *P. ostreatus* fruiting bodies.

Fig. 6. Percent of DPPH radical scavenging activity at different concentrations of β -glucans compared with the universal reducing power (Vitamin C) showing means and standard deviation error bars from duplicate samples.

bonds. The results revealed the presence of an NH group of frequency 3344 cm^{-1} , =C–H group at the frequency of 3078 cm^{-1} , C–H group at 2899 cm^{-1} , also the presence of C=O, N–O, C–H and C–N groups at the frequencies of (1685 , 1564 , 1413 and 1066 cm^{-1}) respectively. However, the frequencies that are less than 1000 cm^{-1} belong to the heavy elements. The data are shown in Fig. 5 and Table 2.

3.4. Antioxidant activity of β -glucans polysaccharide

As shown in Fig. 6, the percentage values of DPPH radical scavenging activity were increased gradually with an increase of the concentrations of β -glucans and the universal reducing power (vitamin C). The

Fig. 7. Antimicrobial activity of β -glucans against some pathogenic bacteria.

Table 1. The retention times of peaks of a polysaccharide β -glucans.

Reten. Time	Result table (Uncal – F: \ β -glucans (10 ppm) – Detector 1)				Compound Name
	Area [mAU.s]	Height [mAU]	Area [%]	Height W 05 [min]	
1	5.927	3299.544	208.935	100.0	100.0 0.25
Total	3299.544	208.935	100.0	100.0	

Table 2. Result of FTIR spectroscopy frequency ranges and absorptions for functional groups.

S. No	Frequency	Assignment
1	3344	NH, OH
2	3078	= C–H
3	2899	C–H stretch
4	1685	C=O
5	1564	N–O, C–C
6	1519	C–C
7	1413	C–H
8	1348	N–O
9	1066	C–N
10	819	C–CL
11	657	C–Br

result also revealed that β -glucans exhibit more percent of scavenging activity (8.59, 12.36, 18.56, 23.69, 44.66, and 80.36%) than vitamin C (2.45, 4.75, 11.95, 19.45, 36.49, and 72.46%) at the concentrations of 31.2, 64.4, 125, 250, 500 and 800 μ g/mL, respectively.

Moreover, the standard deviations with β -glucans were (0.834386002, 0.509116882, 0.791959595, 0.975807358, 0.933380951 and 0.862670273), while the standard deviations with vitamin C were (0.636396103, 0.636396103, 0.777817459, 0.707106781, 0.721248917 and 0.763675324), at the above concentrations, respectively (Fig. 6).

Table 3. The average numbers of antibacterial activity of β -glucans. Inhibition zone (Diameter in mm).

S. No	Microorganisms	Distilled water (-ve control)	Chloramphenicol mg/mL 40 (+ve control)	β -glucans 100 mg/mL
1	<i>P. aeruginosa</i>	0	22	45
2	<i>E. coli</i>	0	21	23
3	<i>S. aureus</i>	0	20	13
4	<i>P. mirabilis</i>	0	24	24

Table 4. Effect of purified β -glucans polysaccharide isolated from *P. ostreatus* on MCF-7 and WRL-68 lines using MTT assay for 24 h at 37 °C.

Cancer cell line	Concentrations (μ g/mL)				
	1.0	1.5	2.0	2.5	3.0
Percentage of growth-inhibiting of cancer cells					
MCF-7	18	24	50	59	62
WRL-68	4	6	9	13	22

3.5. Antimicrobial activity of β -glucans polysaccharide

The result showed remarkable antimicrobial activity against *P. aeruginosa*, where the highest growth inhibition zone was (45 mm) at 100 mg/mL, followed by *P. mirabilis* (24 mm) and *E. coli* (23 mm). The lowest antimicrobial activity of β -glucans appeared against *S. aureus* (13 mm), as shown in Fig. 7 and Table 3.

3.6. Cytotoxic activity of β -glucans polysaccharide

The cytotoxicity results of β -glucans on MCF-7 and WRL-68 lines showed a variation in the cytotoxicity effect at all concentrations of purified β -glucans polysaccharide isolated from *P. ostreatus*. Remarkably, the MCF-7 line was more affected in increasing the percentage of growth-inhibiting than the WRL-68 line. Significantly, it was found that the cytotoxicity increases with increasing the glucose concentration, which is reflected in increasing the percentage of inhibition of cancer cells. The highest values were 62% and 22% at a concentration of 3.0 μ g/ml in MCF-7 and WRL-68 lines, respectively. While the lowest values of the percentage of growth inhibition were 18% and 4% at a concentration of 1.0 μ g/ml in MCF-7 and WRL-68 lines, respectively (Table 4).

4. Discussion

The 657 bp fragment of ITS region (GenBank accession no. MW457626) of fruiting bodies isolated from wild *P. ostreatus* was confirmed (Fig. 2), and the homology searches were blasted against the NCBI database (Fig. 3). This region of nuclear ribosomal DNA has been widely used (as a common sequenced DNA region) for molecular genetic identification in various fungi [31]. The FTIR spectrum result revealed the presence of β -glycosidic bonds such as the presence of (NH, C–H, =C–H, C=O, and other groups) (Fig. 5 and Table 2). FTIR spectroscopy is considered an outstanding technique for structural analysis of polysaccharides [58,59]. This procedure can easily predict the anomeric arrangement and position of the glycosidic bonds in glucans. It detects the glucans in different raw materials especially in the crude with high molecular fractions. For instance, two essential kinds of glucans can be found in the fruiting bodies of mushrooms which are linear (1 $\rightarrow$ 3)- α -D and branched (1 $\rightarrow$ 3) (1 $\rightarrow$ 6)- β -D glucans [60]. Reactive oxygen species are necessary to initiate mutagenesis. The inhibitory effect resulting from antioxidant

activities may lead to DNA mutagenesis produced by oxidative stress [61]. Also, oxidative stress has been considered to be one of the primary causal factors for different diseases and ageing, the antioxidant capacity of β -glucans was examined. The DPPH radical scavenging properties can be attributed to their hydrogen donating capacity. Our findings (Fig. 6) are consistent with previous results [62] who mentioned that antioxidant efficiencies by inhibitory concentration on DPPH showed a significant difference compared to universal antioxidants such as ascorbic acid. Moreover, the current result is agreed with [63] findings who reported that exopolysaccharides extracts from *G. lucidum* at 50, 100, 250, 500 and 1000 μ g/ml revealed maximum antioxidant ability ($82.30 \pm 1.2\%$).

As natural antimicrobial compounds are important, mushrooms need these compounds as a source of antibiotics during their development. Industrial antibiotics and antimicrobial drugs perhaps are dangerous to human health and often can cause antibiotic resistance [64]. Therefore, using natural antimicrobial compounds is important [65]. Interestingly, β -glucans in fungal cell walls have been detected to have an antimicrobial impact [66]. Our results (Fig. 7 and Table 3) share similarities with [67] findings who mentioned that polysaccharides isolated from *Ganoderma* species have antibacterial effects against different types of bacteria *in vitro*. The current result is also concurred well with [6,18,62,66] who found that mushrooms have antimicrobial agents.

Cancers are mainly caused by the mutations in genetic material as a result of the changes that occur in the sequence of DNA, and these mutations can influence the structure as well as the function of the encoded proteins, then in genetic characteristics. It is found that 90% of the major reasons of cancers are correlated with mutations as a consequence of environmental factors for instance chemicals, radiation, metals, and others [68]. According to our results, it can reasonably assume that β -glucans polysaccharide helps to prevent or decrease the growth of cancer cells and maybe prevent the mutations as well. Our results (Table 4) are consistent with [48] who found that *Ganoderma lucidum* polysaccharide (GLP) isolated from *G. lucidum* which is the major compound of β -glucans can inhibit the mutations, also it is in complete agreement with [69] that β -glucans extracted from oyster mushrooms displays anti-neoplastic activity against MCF-7 line [70]. The reason for the anticancer properties is probably because of the activity of β -glucans polysaccharide, particularly in β -1,3 bound sites [71,72]. The current results are also in agreement with

[18,73] who mentioned that bioactive molecule in *Pleurotus* sp. (oyster mushroom) involves high molecular weight compounds (polysaccharides) especially β -glucans which has anti-cancer effects.

5. Conclusion

To sum up, the ITS region of *P. ostreatus* was molecularly confirmed. Multiple sequence alignment and phylogenetic tree of studied ITS were performed depending on some available sequences in GenBank. The analysis of HPLC and FTIR has revealed the presence of β -glucans polysaccharide in isolated specimens. The current result revealed the highest antioxidant activity of β -glucans was 80.36% at 800 μ g/ml. Moreover, a larger zone of inhibition of bacterial growth was (45 mm) against *P. aeruginosa* at 100 mg/ml. The highest percentage of growth inhibition in MCF-7 line was 62% at 3.0 μ g/ml comparing with 22% in WRL-68 line. It seems that β -glucans polysaccharide of *P. ostreatus* has various health benefits and therapeutic properties; and it can be considered safe alternatives for human food and health.

Acknowledgement

This work was performed at the research unit of the Department of the Biology/University of Mosul. The authors would like to thank the Research Center of the Biotechnology at the University of Al-Nahrain for their kind assistance to provide MCF-7 and WRL-68 cancer cell lines.

References

- [1] S.P. Wasser, A.L. Weis, Medicinal properties of substances occurring in higher basidiomycetes mushrooms: current perspectives, *Int. J. Med. Mushrooms* 1 (1999) 31–62, <https://doi.org/10.1615/intjmedmushrooms.v1.i1.30>.
- [2] P. Bobek, S. Galbvy, Effect of pleuran (beta-glucan from *Pleurotus ostreatus*) on the antioxidant status of the organism and on dimethylhydrazine-induced precancerous lesions in rat colon, *Br. J. Biomed. Sci.* 58 (2001) 164–168, <https://search.proquest.com/openview/4308d566b47ec9c20b4b34e5967ed288/1?pq-origsite=gscholar&cbl=4969>.
- [3] E. Guillaumon, A. García-Lafuente, M. Lozano, M.A. Rostagno, A. Villares, J.A. Martínez, Edible mushrooms: role in the prevention of cardiovascular diseases, *Fitoterapia* 81 (2010) 715–723, <https://doi.org/10.1016/j.fitote.2010.06.005>.
- [4] S. Wasser, Medicinal mushroom science: current perspectives, advances, evidences, and challenges, *Biomed. J.* 37 (2014) 345–356, <https://doi.org/10.4103/2319-4170.138318>.
- [5] M.E. Valverde, T. Hernández-Pérez, O. Paredes-López, Edible mushrooms: improving human health and promoting quality life, *Int. J. Microbiol.* 2015 (2015) 1–14, <https://doi.org/10.1155/2015/376387>.
- [6] C. Surekha, C. Kaladhar, D.S.V.G.K. Raju, S.J.R. Haseena, Evaluation of antioxidant and antimicrobial potentiality of some edible mushrooms, *Int. J. Adv. Biotech. Res.* 2 (2011) 130–134, https://www.researchgate.net/publication/213534731_EVALUATION_OF_ANTIOXIDANT_AND_ANTIMICROBIAL_POTENTIALITY_OF_SOME_EDIBLE_MUSHROOMS.
- [7] V. Lavelli, C. Proserpio, F. Gallotti, M. Laureati, E. Pagliarini, Circular reuse of bio-resources: the role of *Pleurotus* spp. in the development of functional foods, *Food Funct.* 9 (2018) 1353–1372, <https://doi.org/10.1039/C7FO01747B>.
- [8] F. Gallotti, V. Lavelli, The effect of UV irradiation on vitamin D2 content and antioxidant and antiglycation activities of mushrooms, *Foods* 9 (2020) 1087, <https://doi.org/10.3390/foods9081087>.
- [9] E. Pereira, L. Barros, A. Martins, I.C. Ferreira, Towards chemical and nutritional inventory of Portuguese wild edible mushrooms in different habitats, *Food Chem.* 130 (2012) 394–403, <https://doi.org/10.1016/j.foodchem.2011.07.057>.
- [10] K.A. Thillaimaharani, K. Sharmila, P. Thangaraju, M. Karthick, M. Kalaiselvam, Studies on antimicrobial and antioxidant properties of oyster mushroom *Pleurotus florida*, *Int. J. Pharm. Sci. Res.* 4 (2013) 1540, <https://ijpsr.com/bft-article/studies-on-antimicrobial-and-antioxidant-properties-of-oyster-mushroom-pleurotus-florida/?view=fulltext>.
- [11] M.A. Khan, M. Tania, Nutritional and medicinal importance of *Pleurotus* mushrooms: an overview, *Food Rev. Int.* 28 (2012) 313–329, <https://doi.org/10.1080/87559129.2011.637267>.
- [12] K. Deepalakshmi, M. Sankaran, *Pleurotus ostreatus*: an oyster mushroom with nutritional and medicinal properties, *J. Biochem. Technol.* 5 (2014) 718–726, <https://jbiochemtech.com/storage/models/article/NG23jvirk6MsPU83nHuA6CbEMW8XcyYx1abn0BuLtqBOKsnuWPknyki9rj5/pleurtus-ostreatus-an-oyster-mushroom-with-nutritional-and-medicinal-properties.pdf>.
- [13] R.C.G. Corrêa, A.H.P. de Souza, R.C. Calhelha, L. Barros, J. Glamoclija, M. Sokovic, R.M. Peralta, A. Bracht, I.C. Ferreira, Bioactive formulations prepared from fruiting bodies and submerged culture mycelia of the Brazilian edible mushroom *Pleurotus ostreatus* Singer, *Food Funct.* 6 (2015) 2155–2164, <https://doi.org/10.1039/C5FO00465A>.
- [14] A. SyNytSyA, K. Mickova, I. Jablonsky, M. SIUKOVÁ, J. Copikova, Mushrooms of genus *Pleurotus* as a source of dietary fibres and glucans for food supplements, *Czech J. Food Sci.* 26 (2008) 441–446, <https://doi.org/10.17221/1361-CJFS>.
- [15] Y. Patel, R. Naraian, V.K. Singh, Medicinal properties of *Pleurotus* species (oyster mushroom): a review, *World J. Fungal Plant Biol.* 3 (2012) 1–12, <https://doi.org/10.5829/idosi.wjfpb.2012.3.1.303>.
- [16] U. Kües, Y. Liu, Fruiting body production in basidiomycetes, *Appl. Microbiol. Biotechnol.* 54 (2000) 141–152, <https://doi.org/10.1007/s002530000396>.
- [17] P. Manzi, L. Gambelli, S. Marconi, V. Vivanti, L. Pizzoferrato, Nutrients in edible mushrooms: an inter-species comparative study, *Food Chem.* 65 (1999) 477–482, [https://doi.org/10.1016/S0308-8146\(98\)00212-X](https://doi.org/10.1016/S0308-8146(98)00212-X).
- [18] H.J. Morris, Y. Beltrán, G. Llauro, P.L. Batista, I. Perraud-Gaime, N. García, S. Moukha, R.C. Bermúdez, P. Cos, E. Hernández, J.C. Díez, Mycelia from *Pleurotus* sp. (oyster mushroom): a new wave of antimicrobials, anticancer and antioxidant bio-ingredients, *Int. J. Phytocosmet. Nat. Ingrid.* 4 (2017) 1–9, <https://doi.org/10.15171/ijpni.2017.03>.
- [19] S. Patel, A. Goyal, Recent developments in mushrooms as anti-cancer therapeutics: a review, *3 Biotech.* 2 (2012) 1–15, <https://doi.org/10.1007/s13205-011-0036-2>.
- [20] S. Rahar, G. Swami, N. Nagpal, M.A. Nagpal, G.S. Singh, Preparation, characterization, and biological properties of β -glucans, *J. Adv. Pharm. Technol. Res.* 2 (2011) 94, <https://doi.org/10.4103/F2231-4040.82953>.
- [21] X. Pang, Z. Chen, X. Gao, W. Liu, M. Slavin, W. Yao, L.L. Yu, Potential of a novel polysaccharide preparation (GLPP) from Anhui-Grown *Ganoderma lucidum* in tumor treatment and immunostimulation, *J. Food Sci.* 72 (2007) S435–S442, <https://doi.org/10.1111/j.1750-3841.2007.00431.x>.

- [22] W. Xu, F. Zhang, Y. Luo, L. Ma, X. Kou, K. Huang, Antioxidant activity of a water-soluble polysaccharide purified from *Pteridium aquilinum*, Carbohydr. Res. 344 (2009) 217–222, <https://doi.org/10.1016/j.carres.2008.10.021>.
- [23] J.H. Cho, J.Y. Lee, M.G. Lee, H.N. Oh, D.H. Kang, C.S. Jhune, Comparative analysis of useful β -glucan and polyphenol in the fruiting bodies of *Ganoderma* spp., J. Mushrooms 11 (2013) 164–170, <https://doi.org/10.14480/JM.2013.11.3.164>.
- [24] M. Kohguchi, T. Kunikata, H. Watanabe, N. Kudo, T. Shibuya, T. Ishihara, K. Iwaki, M. Ikeda, S. Fukuda, M. Kurimoto, Immuno-potentiating effects of the antler-shaped fruiting body of *Ganoderma lucidum* (Rokkaku-Reishi), Biosci. Biotechnol. Biochem. 68 (2004) 881–887, <https://doi.org/10.1271/bbb.68.881>.
- [25] J.H. Park, Y.M. Jin, S. Hwang, D.H. Cho, D.H. Kang, I. Jo, Uric acid attenuates nitric oxide production by decreasing the interaction between endothelial nitric oxide synthase and calmodulin in human umbilical vein endothelial cells: a mechanism for uric acid-induced cardiovascular disease development, Nitric Oxide 32 (2013) 36–42, <https://doi.org/10.1016/j.niox.2013.04.003>.
- [26] O. Rop, J. Mlcek, T. Jurikova, Beta-glucans in higher fungi and their health effects, Nutr Rev 67 (2009) 624–631, <https://doi.org/10.1111/j.1753-4887.2009.00230.x>.
- [27] M. Sari, A. Prange, J.I. Lelley, R. Hambitzer, Screening of beta-glucan contents in commercially cultivated and wild growing mushrooms, Food Chem. 216 (2017) 45–51, <https://doi.org/10.1016/j.foodchem.2016.08.010>.
- [28] J. Nitschke, H. Modick, E. Busch, R.W. Von Rekowski, H.J. Altenbach, H. Mölleken, A new colorimetric method to quantify β -1, 3-1, 6-glucans in comparison with total β -1, 3-glucans in edible mushrooms, Food Chem. 127 (2011) 791–796, <https://doi.org/10.1016/j.foodchem.2010.12.149>.
- [29] J. Liu, Y. Sun, H. Yu, C. Zhang, L. Yue, X. Yang, L. Wang, J. Liu, Purification and identification of one glucan from golden oyster mushroom (*Pleurotus citrinopileatus* (Fr.) Singer), Carbohydr. Polym. 87 (2012) 348–352, <https://doi.org/10.1016/j.carbpol.2011.07.059>.
- [30] K.G. Peay, P.G. Kennedy, T.D. Bruns, Fungal community ecology: a hybrid beast with a molecular master, Bioscience 58 (2008) 799–810, <https://doi.org/10.1641/B580907>.
- [31] S.Y. Abdulhadi, R.N. Gergees, G.Q. Hasan, Molecular identification, antioxidant efficacy of phenolic compounds, and antimicrobial activity of beta-carotene isolated from fruiting bodies of *Suillus* sp., Karbala Int. J. Mod. Sci. 6 (2020) 365–374, <https://doi.org/10.33640/2405-609X.1966>.
- [32] C.L. Schoch, K.A. Seifert, S. Huhndorf, V. Robert, J.L. Spouge, C.A. Levesque, W. Chen, Fungal Barcoding Consortium, Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi, Proc. Natl. Acad. Sci. 109 (2012) 6241–6246, <https://doi.org/10.1073/pnas.1117018109>.
- [33] I.S.E. Ueitele, P. Chimwamurombe, P.N. Kadhila-Muandingi, Optimization of indigenous *Ganoderma lucidum* productivity under cultivation in Namibia, University of Namibia, Int. Sci. Technol. J. Namibia 3 (2014) 35–41, https://www.researchgate.net/publication/280011621_Optimization_of_indigenous_Ganoderma_lucidum_productivity_under_cultivation_in_Namibia.
- [34] D.L. Largent, How to identify mushrooms to genus I: macroscopic Features, Mad River Press, Eureka, 1977, <https://dokumen.pub/how-to-identify-mushrooms-to-genus-i-macroscopic-features-revised-edition-0916422003.html>.
- [35] D.L. Largent, D. Johnson, R. Watling, How to identify mushrooms to genus III: microscopic Features, Mad River Press, Eureka, 1977, <http://files.shroomery.org/attachments/21366346-Optimized/HowToIdentifyMushrooms/To/Genus/III/Microscopic/Features/-/OCR.pdf>.
- [36] R. Singer, Fungi of northern Brazil, Universidade do Recife. Instituto de Micologia, 1961.
- [37] R. Singer, The agaricales in modern taxonomy, Koeltz Scientific Books, Koenigstein, Germany, 1986.
- [38] D.N. Pegler, *Pleurotus* (agaricales) in India, Nepal and Pakistan, Kew. Bull. 31 (1977) 501–510, <https://doi.org/10.2307/4119394>.
- [39] D.N. Pegler, A preliminary agaric flora of East Africa, Her Majesty's Stationery Office, London, UK, 1977, <https://www.cabdirect.org/cabdirect/abstract/19780644719>.
- [40] D.N. Pegler, Agarics of são paulo, Royal Botanic Gardens, Kew, 1997.
- [41] E.J.H. Corner, The agaric genera *Lentinus*, *Panus*, and *Pleurotus* with particular reference to Malaysian species, Beih Nova Hedwigia 69 (1981) 1–169, <http://pascal-francis.inist.fr/vibad/index.php?action=getRecordDetail&idt=PASCAL82X0100563>.
- [42] O. Hilber, Die gattung *Pleurotus*, Bibl. Mycol. 87 (1982) 1–448, <https://ci.nii.ac.jp/naid/10011769707/>.
- [43] O. Hilber, The genus *Pleurotus* (Fr.) Kummer (2), Selbstverlag, Kelheim, 1997.
- [44] G. Guzmán, L. Montoya, D. Salmones, V.M. Bandala, Studies of the genus *Pleurotus* (Basidiomycotina), II. P. djamor in Mexico and in other Latin-American Countries, taxonomic confusions, distribution and semi-industrial culture, Crypt. Bot. 3 (1993) 213–220, <https://agris.fao.org/agris-search/search.do?recordID=US201301778825>.
- [45] B.E. Lechner, J.E. Wright, E. Albertó, The genus *Pleurotus* in Argentina, Mycologia 96 (2004) 845–858, <https://doi.org/10.1080/15572536.2005.11832931>.
- [46] K. O'Donnell, Phylogenetic evidence indicates the important mycotoxigenic strains Fn-2, Fn-3, Fn-2B and Fn-M represent a new species of *Fusarium*, JSM Mycotoxins 45 (1997) 1–10, https://doi.org/10.2520/myco1975.1997.45_1.
- [47] M.Y. Lung, J.C. Tsai, Antioxidant properties of polysaccharides from the Willow Bracket medicinal mushroom, *Phellinus igniarius* (L.) Quél. (Aphyllphoromycetideae) in submerged culture, Int. J. Med. Mushrooms 11 (2009) 383–394, <https://doi.org/10.1615/IntJMedMushr.v11.i4.50>.
- [48] H.R. Lee, H.B. Lim, Antimutagenic and antioxidative effects of polysaccharides isolated from the water extract of *Ganoderma lucidum*, J. Appl. Pharm. Sci. 9 (2019) 1–7, <https://doi.org/10.7324/JAPS.2019.90401>.
- [49] S. Sumathi, G.T. Iswariya, B. Sivaprabha, B. Dharani, P. Radha, P.R. Padma, Comparative study of radical scavenging activity and phytochemical analysis of fresh and dry rhizomes of *Curcuma zedoaria*, Int. J. Pharm. Sci. Res. 4 (2013) 1069, <https://citeseeerx.ist.psu.edu/viewdoc/download?doi=10.1.1.300.4810&rep=rep1&type=pdf>.
- [50] C. Perez, Antibiotic assay by agar-well diffusion method, Acta Biol. Med. Exp. 15 (1990) 113–115.
- [51] S. Visalakchi, J. Muthumary, Antimicrobial activity of the new endophytic *Monodictys castaneae* SVJM139 pigment and its optimization, Afr. J. Microbiol. Res. 3 (2009) 550–556, <https://doi.org/10.5897/AJMR.9000078>.
- [52] M. Osínska-Jaroszuk, M. Jaszek, M. Mizerska-Dudka, A. Błachowicz, T.P. Rejczak, G. Janusz, J. Wydrych, J. Polak, A. Jarosz-Wilkolazka, M. Kandefer-Szerszeń, Exopolysaccharide from *Ganoderma applanatum* as a promising bioactive compound with cytostatic and antibacterial properties, Biomed. Res. Int. 2014 (2014) 1–10, <https://doi.org/10.1155/2014/743812>.
- [53] P.H. Sneath, R.R. Sokal, Numerical taxonomy. The principles and practice of numerical classification, 1973, <https://www.cabdirect.org/cabdirect/abstract/19730310919>.
- [54] J. Felsenstein, Confidence limits on phylogenies: an approach using the bootstrap, Evolution 39 (1985) 783–791, <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>.
- [55] K. Tamura, M. Nei, S. Kumar, Prospects for inferring very large phylogenies by using the neighbor-joining method, Proc. Natl. Acad. Sci. 101 (2004) 11030–11035, <https://doi.org/10.1073/pnas.0404206101>.
- [56] S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, Mega X: molecular evolutionary genetics analysis across computing platforms, Mol. Biol. Evol. 35 (2018) 1547–1549, <https://doi.org/10.1093/molbev/msy096>.

- [57] G. Mradu, S. Saumyakanti, M. Sohini, M. Arup, HPLC profiles of standard phenolic compounds present in medicinal plants, *Int. J. Pharmacogn. Phytochem. Res.* 4 (2012) 162–167. <http://www.ijppr.com/>.
- [58] B.M. Prado, S. Kim, B.F. Özen, L.J. Mauer, Differentiation of carbohydrate gums and mixtures using Fourier transform infrared spectroscopy and chemometrics, *J. Agric. Food Chem.* 53 (2005) 2823–2829, <https://doi.org/10.1021/jf0485537>.
- [59] M. Kacurakova, P. Capek, V. Sasinkova, N. Wellner, A. Ebringerova, FT-IR study of plant cell wall model compounds: pectic polysaccharides and hemicelluloses, *Carbohydr. Polym.* 43 (2000) 195–203, [https://doi.org/10.1016/S0144-8617\(00\)00151-X](https://doi.org/10.1016/S0144-8617(00)00151-X).
- [60] A. Synytsya, M. Novak, Structural analysis of glucans, *Ann. Transl. Med.* 2 (2014) 1–14, <https://doi.org/10.3978/Fj.issn.2305-5839.2014.02.07>.
- [61] N.R. Jena, DNA damage by reactive species: mechanisms, mutation and repair, *J. Biosci.* 37 (2012) 503–517, <https://doi.org/10.1007/s12038-012-9218-2>.
- [62] M.M.H. Chowdhury, K. Kubra, S.R. Ahmed, Screening of antimicrobial, antioxidant properties and bioactive compounds of some edible mushrooms cultivated in Bangladesh, *Ann. Clin. Microbiol. Antimicrob.* 14 (2015) 1–6, <https://doi.org/10.1186/s12941-015-0067-3>.
- [63] S. Mahendran, K.T.K. Anandapandian, T. Shankar, C. Chellaram, P. Vijayabaskar, Antioxidant properties of *Ganoderma lucidum* crude exopolysaccharide, *Indian J. Innov. Dev.* 1 (2012) 1–6.
- [64] Ö. Özcan, F. Ertan, Beta-glucan content, antioxidant and antimicrobial activities of some edible mushroom species, *Food Sci. Technol.* 6 (2018) 47–55, <https://doi.org/10.13189/fst.2018.060201>.
- [65] S.Y. Abdulhadi, G.Q. Hasan, R.N. Gergees, Molecular detection and antimicrobial activity of Endophytic fungi isolated from a medical plant *Rosmarinus officinalis*, *Ann. Trop. Med. Publ. Health* 23 (2020) 231–384, <https://doi.org/10.36295/ASRO.2020.231384>.
- [66] M. Akyuz, A. Onganer, P. Erecevit, S. Kirbag, Antimicrobial activity of some edible mushrooms in the eastern and southeast Anatolia region of Turkey, *GU J. Sci.* 23 (2010) 125–130. <https://dergipark.org.tr/en/pub/gujs/issue/7386/96882>.
- [67] Y. Gao, W. Tang, H. Gao, E. Chan, J. Lan, X. Li, S. Zhou, Antimicrobial activity of the medicinal mushroom *Ganoderma*, *Food Rev. Int.* 21 (2005) 211–229, <https://doi.org/10.1081/FRI-200051893>.
- [68] P. Anand, S.G. Thomas, A.B. Kunnumakkara, C. Sundaram, K.B. Harikumar, B. Sung, S.T. Tharakan, K. Misra, I.K. Priyadarsini, K.N. Rajasekharan, Biological activities of curcumin and its analogues (Congeners) made by man and Mother Nature, *Biochem. Pharmacol.* 76 (2008) 1590–1611, <https://doi.org/10.1016/j.bcp.2008.08.008>.
- [69] K.R. Martin, S.K. Brophy, Commonly consumed and specially dietary mushrooms reduce cellular proliferation in MCF-7 human breast cancer cells, *Exp. Biol. Med.* 235 (2010) 1306–1314, <https://doi.org/10.1258/Febr.2010.010113>.
- [70] Josef Augustin, Grażyna Jaworska, Alexander Dandar, Kamil Cejpek, Bocznik ostrogowy [Pleurotus ostreatus] jako źródło beta-D-glukanów, *Zywność Nauka Technologia Jakość* 14 (2007) 170–176. <http://agro.icm.edu.pl/agro/element/bwmeta1.element.agro-article-232297a3-e071-46c7-b5ca-71fb3e2a1867>.
- [71] J. Rajewska, B. Bałasińska, Związki biologicznie aktywne zawarte w grzybach jadalnych i ich korzystny wpływ na zdrowie [Biologically active compounds of edible mushrooms and their beneficial impact on health], *Post. Hig. Med. Dośw.* 58 (2004) 352–357. PMID: 15536392.
- [72] I.B. Jeff, E. Fan, M. Tian, C. Song, J. Yan, Y. Zhou, In vivo anticancer and immunomodulating activities of mannogalactoglucan-type polysaccharides from *Lentinus edodes* (Berkeley) Singer, *Cent. Eur. J. Immunol.* 41 (2016) 47–53, <https://doi.org/10.5114/Fceji.2015.56962>.
- [73] D.D. De Silva, S. Rapior, E. Sudarman, M. Stadler, J. Xu, S.A. Alias, K.D. Hyde, Bioactive metabolites from macrofungi: ethnopharmacology, biological activities and chemistry, *Fungal Divers* 62 (2013) 1–40, <https://doi.org/10.1007/s13225-013-0265-2>.